



“Paxman made this very difficult journey better for me.”

Brenda.

Hair loss is the most feared and common side effect of chemotherapy.

Scalp Cooling can be used to minimize chemotherapy-induced alopecia and can positively impact the confidence, sense of control and hope of patients who use it.

Paxman is the global leader in scalp cooling, used by the main Cancer Centers across the USA.


View our clinical efficacy data and find out how you can give your patients back some control visit:

www.paxmanUSA.com

Offer your patients the option of scalp cooling, they will...

THANK YOU

PAXMAN^o
CHANGING THE FACE OF CANCER




PAXMAN^o
CHANGING THE FACE OF CANCER

HCP@paxmanUSA.com
www.paxmanUSA.com



ISSUE 1 0719



NCCN
RECOMMENDED

NCCN Clinical Practice Guidelines In Oncology (NCCN Guidelines[®]) now recommend scalp cooling as a category 2A treatment option for patients with invasive breast cancer. To reduce incidence of chemotherapy-induced alopecia in those receiving chemotherapy.

Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Breast Cancer V.1.2019. © National Comprehensive Cancer Network, Inc. 2019. All rights reserved. Accessed March 14, 2019. To view the most recent and complete version of the guideline, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

PAXMAN^o
CHANGING THE FACE OF CANCER

Invasive Breast Cancer

PREOPERATIVE/ADJUVANT THERAPY REGIMENS^{a,b,c,d,e,f}HER2-Negative^g

Preferred regimens:

- Dose-dense AC (doxorubicin/cyclophosphamide) followed by paclitaxel every 2 weeks^h
- Dose-dense AC (doxorubicin/cyclophosphamide) followed by weekly paclitaxelⁱ
- TC (docetaxel and cyclophosphamide)
- If triple-negative breast cancer and residual disease after preoperative therapy with taxane-, alkylator-, and anthracycline- based chemotherapy: capecitabine^j

Useful in certain circumstances:

- Dose-dense AC (doxorubicin/cyclophosphamide)
- AC (doxorubicin/cyclophosphamide) every 3 weeks (category 2B)
- CMF (cyclophosphamide/methotrexate/fluorouracil)
- AC followed by weekly paclitaxel

Other recommended regimens:

- AC followed by docetaxel every 3 weeks
- EC (epirubicin/cyclophosphamide)
- TAC (docetaxel/doxorubicin/cyclophosphamide)

^a Retrospective evidence suggests that anthracycline-based chemotherapy regimens may be superior to non-anthracycline-based regimens in patients with HER2-positive tumors.

^b Randomized clinical trials demonstrate that the addition of a taxane to anthracycline-based chemotherapy provides an improved outcome.

^c CMF and radiation therapy may be given concurrently, or the CMF may be given first. All other chemotherapy regimens should be given prior to radiotherapy.

^d Chemotherapy and endocrine therapy used as adjuvant therapy should be given sequentially with endocrine therapy following chemotherapy.

^e Nab-paclitaxel may be substituted for paclitaxel or docetaxel due to medical necessity (ie, hypersensitivity reaction). If substituted for weekly paclitaxel or docetaxel, then the weekly dose of nab-paclitaxel should not exceed 125mg/m².

^f **Consider scalp cooling to reduce the incidence of chemotherapy-induced alopecia for patients receiving neoadjuvant/adjuvant chemotherapy. Results may be less effective with anthracycline-containing regimens.**

^g The regimens listed for HER-2 negative are all category 1 (except where indicated) when used in the adjuvant setting.

^h It would be acceptable to change the administration sequence to paclitaxel followed by dose-dense AC.

ⁱ Capecitabine 1,000 – 1250 mg/m² PO twice daily on days 1 -14. Cycled every 21 days for 6 – 8 cycles. Masuda N, Lee SJ, Ohtani S, et al. Adjuvant capecitabine for breast cancer after preoperative chemotherapy. N Engl J Med 2017;376:2147-2159.

Invasive Breast Cancer

PREOPERATIVE/ADJUVANT THERAPY REGIMENS^{a,b,c,d,e,f}HER2-Positive^j

Preferred regimens:

- AC followed by T + trastuzumab^k
- (doxorubicin/cyclophosphamide followed by paclitaxel plus trastuzumab, various schedules)
- AC followed by T + trastuzumab + pertuzumab^l (doxorubicin/cyclophosphamide followed by paclitaxel plus trastuzumab plus pertuzumab)
- Paclitaxel + trastuzumab
- TCH (docetaxel/carboplatin/ trastuzumab)
- TCH (docetaxel/carboplatin/ trastuzumab) + pertuzumab
- If no residual disease after preoperative therapy or no preoperative therapy: Complete up to one year of HER2-targeted therapy with trastuzumab (category 1) ± pertuzumab.^m
- If residual disease after preoperative therapy: Ado- trastuzumab emtansine (category 1) alone n If ado- trastuzumab emtansine discontinued for toxicity, then trastuzumab (category 1) ± pertuzumab to complete one year of therapy.^m

Useful in certain circumstances:

- Docetaxel + cyclophosphamide + trastuzumabⁿ

Other recommended regimens:

- AC followed by docetaxel + trastuzumab^o (doxorubicin/cyclophosphamide followed by docetaxel plus trastuzumab)
- AC followed by docetaxel + trastuzumab + pertuzumab^o
- (doxorubicin/cyclophosphamide followed by docetaxel plus trastuzumab plus pertuzumab)

^a Retrospective evidence suggests that anthracycline-based chemotherapy regimens may be superior to non-anthracycline-based regimens in patients with HER2-positive tumors.

^b Randomized clinical trials demonstrate that the addition of a taxane to anthracycline-based chemotherapy provides an improved outcome.

^c CMF and radiation therapy may be given concurrently, or the CMF may be given first. All other chemotherapy regimens should be given prior to radiotherapy.

^d Chemotherapy and endocrine therapy used as adjuvant therapy should be given sequentially with endocrine therapy following chemotherapy.

^e Albumin-bound paclitaxel may be substituted for paclitaxel or docetaxel due to medical necessity (ie hypersensitivity reaction). If substituted for weekly paclitaxel or docetaxel, then the weekly dose of nab-paclitaxel should not exceed 125mg/m².

^f **Consider scalp cooling to reduce the incidence of chemotherapy-induced alopecia for patients receiving chemotherapy. Results may be less effective with anthracycline-containing regimens.**

^j Trastuzumab and hyaluronidase-oysk injection for subcutaneous use may be substituted for trastuzumab. It has different dosage and administration instructions compared to intravenous trastuzumab. Do not substitute trastuzumab and hyaluronidase-oysk for or with ado- trastuzumab emtansine.

^k Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided.

^l Paclitaxel + trastuzumab may be considered for patients with low-risk T1,N0,M0, HER2-positive disease, particularly those not eligible for other standard adjuvant regimens due to comorbidities.

^m Consider extended adjuvant neratinib following adjuvant trastuzumab-containing therapy for patients with HR-positive, HER2-positive disease with a perceived high risk or recurrence. The benefit or toxicities associated with extended neratinib in patients who have received pertuzumab or ado-trastuzumab emtansine is unknown.

ⁿ Ado-trastuzumab emtansine 3.6mg/kg cycled every 21 days for 14 cycles. Von Minckwitz G, Huang C, Mano M, et al. Trastuzumab emtansine for residual invasive HER2-positive breast cancer. N Engl J Med 2019 ; 380 :617-628.

Invasive Breast Cancer

CHEMOTHERAPY REGIMENS FOR RECURRENT OR STAGE IV (M1) DISEASE^{a,b}

HER2-Negative

Preferred regimens:

- Anthracyclines
 - Doxorubicin
 - Liposomal doxorubicin
- Taxanes
 - Paclitaxel
- Anti-metabolites
 - Capecitabine
 - Gemcitabine
- Microtubule inhibitors
 - Vinorelbine
 - Eribulin
- PARP inhibitors (options for patients with HER2-negative tumors and germline BRCA1/2 mutation)^c
 - Olaparib^d (category 1)
 - Talazoparib^d (category 1)
- Platinum (option for patients with triple-negative tumors and germline BRCA ½ mutation)^e
 - Carboplatin
 - Cisplatin
- Atezolizumab + albumin-bound paclitaxel (option for patients with PD-L1-positive TNBC)^f

Other recommended regimens^g:

- Cyclophosphamide
- Docetaxel
- Albumin-bound paclitaxel
- Epirubicin
- Ixabepilone

Useful in certain circumstances^h:

- AC (doxorubicin/cyclophosphamide)
- EC (epirubicin/cyclophosphamide)
- CMF (cyclophosphamide/methotrexate/fluorouracil)
- Docetaxel/capecitabine
- GT (gemcitabine/docetaxel)
- Gemcitabine/carboplatin
- Paclitaxel/bevacizumabⁱ

HER2-Positive^g

Preferred regimens:

- Pertuzumab + trastuzumab + docetaxel (category 1)^h
- Pertuzumab + trastuzumab + paclitaxelⁱ

Other recommended regimens^j:

- Ado-trastuzumab emtansine (T-DM1)
- Trastuzumab + paclitaxel ± carboplatin
- Trastuzumab + docetaxel^h
- Trastuzumab + vinorelbine^h
- Trastuzumab + capecitabine
- Lapatinib + capecitabine
- Trastuzumab + lapatinib (without cytotoxic therapy)
- Trastuzumab + other agents^{h,i}

^a Albumin-bound paclitaxel may be substituted for paclitaxel or docetaxel due to medical necessity (ie hypersensitivity reaction). If substituted for weekly paclitaxel or docetaxel, then the weekly dose of nab-paclitaxel should not exceed 125mg/m².

^b **Consider scalp cooling to reduce the incidence of chemotherapy-induced alopecia for patients receiving chemotherapy. Results may be less effective with anthracycline-containing regimens.**

^c Sequential single agents are preferred, but chemotherapy combinations may be used in select patients with high tumor burden, rapidly progressing disease, and visceral crisis.

^d Patients with HER2-negative disease, strongly consider for germline BRCA 1/2 testing.

^e Patients with TNBC, assess PD-L1 biomarker status on tumor-infiltrating immune cells to identify patients most likely to benefit from atezolizumab plus albumin-bound paclitaxel.

^f Randomized clinical trials in metastatic breast cancer document that the addition of bevacizumab to some first- or second-line chemotherapy agents modestly improves time to progression and response rates but does not improve overall survival. The time-to-progression impact may vary among cytotoxic agents and appears greatest with bevacizumab in combination with weekly paclitaxel.

^g Trastuzumab and hyaluronidase-oysk injection for subcutaneous use may be substituted for trastuzumab. It has different dosage and administration instructions compared to intravenous trastuzumab. Do not substitute trastuzumab and hyaluronidase-oysk for or with ado-trastuzumab emtansine.

^h Patients previously treated with chemotherapy plus trastuzumab in the absence of pertuzumab in the metastatic setting may be considered for one line of therapy including both trastuzumab plus pertuzumab in combination with or without cytotoxic therapy (such as vinorelbine or taxane). Further research is needed to determine the ideal sequencing strategy for anti-HER2 therapy.

ⁱ Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided.

^j Trastuzumab may be safely combined with all non-anthracycline containing preferred and other single agents listed above for recurrent or metastatic breast cancer.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical, trial. Participation in clinical trials is especially encouraged.

Adapted with permission from the for Breast Cancer version 1.2019 - March 14 2019 Version 1.2019, 03/14/19 © 2019 National Comprehensive Cancer Network® (NCCN®). All rights reserved. NCCN Guidelines® and this illustration may not be reproduced in any form without express written permission of NCCN. To view the most recent and complete version of the NCCN Guidelines go online to NCCN.org. The NCCN Guidelines are a work in progress that may be refined as often as new significant data becomes available. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical, trial. Participation in clinical trials is especially encouraged.

Adapted with permission from the for Breast Cancer version 1.2019 - March 14 2019 Version 1.2019, 03/14/19 © 2019 National Comprehensive Cancer Network® (NCCN®). All rights reserved. NCCN Guidelines® and this illustration may not be reproduced in any form without express written permission of NCCN. To view the most recent and complete version of the NCCN Guidelines go online to NCCN.org. The NCCN Guidelines are a work in progress that may be refined as often as new significant data becomes available. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical, trial. Participation in clinical trials is especially encouraged.

Adapted with permission from the for Breast Cancer version 1.2019 - March 14 2019 Version 1.2019, 03/14/19 © 2019 National Comprehensive Cancer Network® (NCCN®). All rights reserved. NCCN Guidelines® and this illustration may not be reproduced in any form without express written permission of NCCN. To view the most recent and complete version of the NCCN Guidelines go online to NCCN.org. The NCCN Guidelines are a work in progress that may be refined as often as new significant data becomes available. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.